

3R Wrist Positioning – Its Geometric Background

A. Gferrer P.J. Zsombor-Murray

Institut f. Geometrie, TU-Graz
Department of Mechanical Engineering, McGill University

Computational Kinematics 2009 - May 6 - 8

Outline

- 1 Overview & the Conventional Approach
- 2 A Slightly Different Approach
- 3 Removing Absolute Conic $C_\infty = x^2 + y^2 + z^2$ from Cyclid
- 4 Conclusion
- 5 If Time Remains ...

Overview

- Define the wrist positioning problem (WP) in terms of given wrist centre point coordinates and Denavit-Hartenberg parameters of a 3R serial chain.

Overview

- Define the wrist positioning problem (WP) in terms of given wrist centre point coordinates and Denavit-Hartenberg parameters of a 3R serial chain.
- Set up the direct kinematics problem and invert it, producing three scalar equations in joint angles u_1, u_2, u_3 .

Overview

- Define the wrist positioning problem (WP) in terms of given wrist centre point coordinates and Denavit-Hartenberg parameters of a 3R serial chain.
- Set up the direct kinematics problem and invert it, producing three scalar equations in joint angles u_1, u_2, u_3 .
- Eliminate the middle joint angle variable, u_2 .

Overview

- Define the wrist positioning problem (WP) in terms of given wrist centre point coordinates and Denavit-Hartenberg parameters of a 3R serial chain.
- Set up the direct kinematics problem and invert it, producing three scalar equations in joint angles u_1, u_2, u_3 .
- Eliminate the middle joint angle variable, u_2 .
- Produce system of two equations linear in $\cos u_1, \sin u_1, \cos u_3, \sin u_3$.

Overview

- Define the wrist positioning problem (WP) in terms of given wrist centre point coordinates and Denavit-Hartenberg parameters of a 3R serial chain.
- Set up the direct kinematics problem and invert it, producing three scalar equations in joint angles u_1, u_2, u_3 .
- Eliminate the middle joint angle variable, u_2 .
- Produce system of two equations linear in $\cos u_1, \sin u_1, \cos u_3, \sin u_3$.
- Make two tangent half angle = t, u substitutions, *i.e.*,

$$\cos(u_1, u_3) = \frac{1-(t,u)^2}{1+(t,u)^2}, \sin(u_1, u_3) = \frac{2(t,u)}{1+(t,u)^2}.$$

Overview

- Define the wrist positioning problem (WP) in terms of given wrist centre point coordinates and Denavit-Hartenberg parameters of a 3R serial chain.
- Set up the direct kinematics problem and invert it, producing three scalar equations in joint angles u_1, u_2, u_3 .
- Eliminate the middle joint angle variable, u_2 .
- Produce system of two equations linear in $\cos u_1, \sin u_1, \cos u_3, \sin u_3$.
- Make two tangent half angle = t, u substitutions, *i.e.*,


$$\cos(u_1, u_3) = \frac{1-(t,u)^2}{1+(t,u)^2}, \sin(u_1, u_3) = \frac{2(t,u)}{1+(t,u)^2}.$$
- Eliminate, say, t and factor out $[2(1 + u^2)]^2$ from octic.

Overview

- Define the wrist positioning problem (WP) in terms of given wrist centre point coordinates and Denavit-Hartenberg parameters of a 3R serial chain.
- Set up the direct kinematics problem and invert it, producing three scalar equations in joint angles u_1, u_2, u_3 .
- Eliminate the middle joint angle variable, u_2 .
- Produce system of two equations linear in $\cos u_1, \sin u_1, \cos u_3, \sin u_3$.
- Make two tangent half angle = t, u substitutions, *i.e.*,


$$\cos(u_1, u_3) = \frac{1-(t,u)^2}{1+(t,u)^2}, \sin(u_1, u_3) = \frac{2(t,u)}{1+(t,u)^2}.$$
- Eliminate, say, t and factor out $[2(1 + u^2)]^2$ from octic.
- Solve for u_3 . Use two equation system to find u_1 and three equation system to find u_2 .

Conventional Layout




- Sequence of vectors \mathbf{a}_i to be transformed,

Conventional Layout




- Sequence of vectors \mathbf{a}_i to be transformed,
- Joint axes g_i , DH parameters a_1, d_2, a_2, b, s

Conventional Layout



- Sequence of vectors \mathbf{a}_i to be transformed,
- Joint axes g_i , DH parameters a_1, d_2, a_2, b, s
- And common normal pedal points N, O, P, Q, K, W

Conventional Layout



- Sequence of vectors \mathbf{a}_i to be transformed,
- Joint axes g_i , DH parameters a_1, d_2, a_2, b, s
- And common normal pedal points N, O, P, Q, K, W

- Consistent with geometric approach. Skew axis angles α_1, α_2 not shown.

Direct Kinematics and Three Scalar Equations

- Rearrangement of closure equation, *i.e.*, direct kinematics to place W if u_i are given

Direct Kinematics and Three Scalar Equations

- Rearrangement of closure equation, *i.e.*, direct kinematics to place W if u_i are given
- $\mathbf{a}_1 + \mathbf{Q}_1 \mathbf{a}_2 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{a}_3 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{Q}_3 \mathbf{a}_4 - \mathbf{w} = \mathbf{0}$

Direct Kinematics and Three Scalar Equations

- Rearrangement of closure equation, *i.e.*, direct kinematics to place W if u_i are given
- $\mathbf{a}_1 + \mathbf{Q}_1 \mathbf{a}_2 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{a}_3 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{Q}_3 \mathbf{a}_4 - \mathbf{w} = \mathbf{0}$
- Yields three scalar equations

Direct Kinematics and Three Scalar Equations

- Rearrangement of closure equation, *i.e.*, direct kinematics to place W if u_i are given
- $\mathbf{a}_1 + \mathbf{Q}_1 \mathbf{a}_2 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{a}_3 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{Q}_3 \mathbf{a}_4 - \mathbf{w} = \mathbf{0}$
- Yields three scalar equations
-

$$q_{i1} a_3 + q_{i2} b_{\mu_3} + q_{i3} b_{\lambda_3} - w_i = 0, \quad i = 1, 2, 3 \quad (1)$$

Direct Kinematics and Three Scalar Equations

- Rearrangement of closure equation, *i.e.*, direct kinematics to place W if u_i are given
- $\mathbf{a}_1 + \mathbf{Q}_1 \mathbf{a}_2 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{a}_3 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{Q}_3 \mathbf{a}_4 - \mathbf{w} = \mathbf{0}$
- Yields three scalar equations



$$q_{i1} a_3 + q_{i2} b \mu_3 + q_{i3} b \lambda_3 - w_i = 0, \quad i = 1, 2, 3 \quad (1)$$

- $q_{ij} \in \mathbf{Q}_i$, u_i = joint angles, $\mu_i = \cos \alpha_i$, $\lambda_i = \sin \alpha_i$.

Direct Kinematics and Three Scalar Equations

- Rearrangement of closure equation, *i.e.*, direct kinematics to place W if u_i are given
- $\mathbf{a}_1 + \mathbf{Q}_1 \mathbf{a}_2 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{a}_3 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{Q}_3 \mathbf{a}_4 - \mathbf{w} = \mathbf{0}$
- Yields three scalar equations
-

$$q_{i1} a_3 + q_{i2} b \mu_3 + q_{i3} b \lambda_3 - w_i = 0, \quad i = 1, 2, 3 \quad (1)$$

- $q_{ij} \in \mathbf{Q}_i$, u_i = joint angles, $\mu_i = \cos \alpha_i$, $\lambda_i = \sin \alpha_i$.

- $\mathbf{Q}_i = \begin{bmatrix} \cos u_i & -\lambda_i \sin u_i & \mu_i \sin u_i \\ \sin u_i & \lambda_i \sin u_i & -\mu_i \cos u_i \\ 0 & \mu_i & \lambda_i \end{bmatrix}, \quad \mathbf{a}_i = \begin{bmatrix} a_i \cos u_i \\ a_i \sin u_i \\ b_i \end{bmatrix}$

Direct Kinematics and Three Scalar Equations

- Rearrangement of closure equation, *i.e.*, direct kinematics to place W if u_i are given
- $\mathbf{a}_1 + \mathbf{Q}_1 \mathbf{a}_2 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{a}_3 + \mathbf{Q}_1 \mathbf{Q}_2 \mathbf{Q}_3 \mathbf{a}_4 - \mathbf{w} = \mathbf{0}$
- Yields three scalar equations
-

$$q_{i1} a_3 + q_{i2} b \mu_3 + q_{i3} b \lambda_3 - w_i = 0, \quad i = 1, 2, 3 \quad (1)$$

- $q_{ij} \in \mathbf{Q}_i$, u_i = joint angles, $\mu_i = \cos \alpha_i$, $\lambda_i = \sin \alpha_i$.

- $\mathbf{Q}_i = \begin{bmatrix} \cos u_i & -\lambda_i \sin u_i & \mu_i \sin u_i \\ \sin u_i & \lambda_i \sin u_i & -\mu_i \cos u_i \\ 0 & \mu_i & \lambda_i \end{bmatrix}, \quad \mathbf{a}_i = \begin{bmatrix} a_i \cos u_i \\ a_i \sin u_i \\ b_i \end{bmatrix}$

- $b_1 = 0, \quad b_2 = d_2 \quad b_3 = b$

Quatric Monomial

- Angle u_2 can be eliminated from the set of three, Eq. 1 to produce a pair in u_i , $i = 1, 3$.

Quatric Monomial

- Angle u_2 can be eliminated from the set of three, Eq. 1 to produce a pair in u_i , $i = 1, 3$.
- $A_i \cos u_1 + B_i \sin u_1 + C_i \cos u_3 + D_i \sin u_3 = 0$

Quatric Monomial

- Angle u_2 can be eliminated from the set of three, Eq. 1 to produce a pair in u_i , $i = 1, 3$.
- $A_i \cos u_1 + B_i \sin u_1 + C_i \cos u_3 + D_i \sin u_3 = 0$
- Tangent half-angle substitutions produce two in t, u after multiplying through with $(1 + t^2)(1 + u^2)$.

Quatric Monomial

- Angle u_2 can be eliminated from the set of three, Eq. 1 to produce a pair in u_i , $i = 1, 3$.
- $A_i \cos u_1 + B_i \sin u_1 + C_i \cos u_3 + D_i \sin u_3 = 0$
- Tangent half-angle substitutions produce two in t, u after multiplying through with $(1 + t^2)(1 + u^2)$.
- $P_i t^2 u^2 + Q_i t^2 u + R_i t u^2 + S_i t^2 + T_i u^2 + U_i t + V_i u + W_i = 0$

Quatric Monomial

- Angle u_2 can be eliminated from the set of three, Eq. 1 to produce a pair in u_i , $i = 1, 3$.
- $A_i \cos u_1 + B_i \sin u_1 + C_i \cos u_3 + D_i \sin u_3 = 0$
- Tangent half-angle substitutions produce two in t, u after multiplying through with $(1 + t^2)(1 + u^2)$.
- $P_i t^2 u^2 + Q_i t^2 u + R_i t u^2 + S_i t^2 + T_i u^2 + U_i t + V_i u + W_i = 0$
- The resultant in t is an octic in u with a removable factor.

Quatric Monomial

- Angle u_2 can be eliminated from the set of three, Eq. 1 to produce a pair in u_i , $i = 1, 3$.
- $A_i \cos u_1 + B_i \sin u_1 + C_i \cos u_3 + D_i \sin u_3 = 0$
- Tangent half-angle substitutions produce two in t, u after multiplying through with $(1 + t^2)(1 + u^2)$.
- $P_i t^2 u^2 + Q_i t^2 u + R_i t u^2 + S_i t^2 + T_i u^2 + U_i t + V_i u + W_i = 0$
- The resultant in t is an octic in u with a removable factor.
- $[2(1 + u^2)]^2 (P u^4 + Q u^3 + R u^2 + S u + T) = 0$ so as to produce up to four real solutions as expected.

Quatric Monomial

- Angle u_2 can be eliminated from the set of three, Eq. 1 to produce a pair in u_i , $i = 1, 3$.
- $A_i \cos u_1 + B_i \sin u_1 + C_i \cos u_3 + D_i \sin u_3 = 0$
- Tangent half-angle substitutions produce two in t, u after multiplying through with $(1 + t^2)(1 + u^2)$.
- $P_i t^2 u^2 + Q_i t^2 u + R_i t u^2 + S_i t^2 + T_i u^2 + U_i t + V_i u + W_i = 0$
- The resultant in t is an octic in u with a removable factor.
- $[2(1 + u^2)]^2 (P u^4 + Q u^3 + R u^2 + S u + T) = 0$ so as to produce up to four real solutions as expected.
- Aside from back-substitution to find angles $u_{1,2}$ the problem is solved **but why is this sort of octic factorable?**

Scenario

- Split 3R triad so g_2 becomes reference frame z -axis.

Scenario

- Split 3R triad so g_2 becomes reference frame z -axis.
- Wrist centre W sweeps a surface of revolution under u_2, u_3 dyad rotation.

Scenario

- Split 3R triad so g_2 becomes reference frame z -axis.
- Wrist centre W sweeps a surface of revolution under u_2, u_3 dyad rotation.
- Under u_3 rotation, meridian circle k is locus of W , centred on K , radius KW , has axis g_3 skew to g_2 .

Scenario

- Split 3R triad so g_2 becomes reference frame z -axis.
- Wrist centre W sweeps a surface of revolution under u_2, u_3 dyad rotation.
- Under u_3 rotation, meridian circle k is locus of W , centred on K , radius KW , has axis g_3 skew to g_2 .
- Cyclid surface Φ generated as k is rotated by joint angle u_2 about axis g_2 .

Scenario


- Split 3R triad so g_2 becomes reference frame z -axis.
- Wrist centre W sweeps a surface of revolution under u_2, u_3 dyad rotation.
- Under u_3 rotation, meridian circle k is locus of W , centred on K , radius KW , has axis g_3 skew to g_2 .
- Cyclid surface Φ generated as k is rotated by joint angle u_2 about axis g_2 .
- Circle l with centre L is locus of wrist target point W_0 as frame with z -axis g_1 *counter-rotates* about reference frame with g_2 as z -axis.

Scenario


- Split 3R triad so g_2 becomes reference frame z -axis.
- Wrist centre W sweeps a surface of revolution under u_2, u_3 dyad rotation.
- Under u_3 rotation, meridian circle k is locus of W , centred on K , radius KW , has axis g_3 skew to g_2 .
- Cyclid surface Φ generated as k is rotated by joint angle u_2 about axis g_2 .
- Circle l with centre L is locus of wrist target point W_0 as frame with z -axis g_1 *counter-rotates* about reference frame with g_2 as z -axis.
- Computational process, reducing a sphero-quartic surface and some questions.

Frame Definitions

- A 3R chain is shown in “home” position wherein



Frame Definitions




- A 3R chain is shown in "home" position wherein
- Common normals NO PQ KW between axes are parallel.

The diagram shows a 3-DOF serial manipulator with three revolute joints. The base is fixed to a ground frame Σ_0 . The first joint is at point O with axis N (vertical). The second joint is at point P with axis Q (along the link OP). The third joint is at point K with axis W (along the link PK). The end effector is a point W . The links are labeled Σ_1 , Σ_2 , and Σ_3 . The Denavit-Hartenberg parameters are listed as follows:

- $NO = a_1$
- $OP = d$
- $PQ = a_2$
- $QK = b$
- $KW = s$


- A 3R chain is shown in “home” position wherein
- Common normals NO PQ KW between axes are parallel.
- Denavit-Hartenberg (DH) parameters are defined.

Frame Definitions




- A 3R chain is shown in "home" position wherein
- Common normals NO PQ KW between axes are parallel.
- Denavit-Hartenberg (DH) parameters are defined.
- Purpose is to place $W \rightarrow W_0$ by

Frame Definitions




- A 3R chain is shown in "home" position wherein
- Common normals NO PQ KW between axes are parallel.
- Denavit-Hartenberg (DH) parameters are defined.
- Purpose is to place $W \rightarrow W_0$ by
- Rotation angles u_1, u_2, u_3 about

Frame Definitions



- A 3R chain is shown in “home” position wherein
- Common normals NO PQ KW between axes are parallel.
- Denavit-Hartenberg (DH) parameters are defined.
- Purpose is to place $W \rightarrow W_0$ by
- Rotation angles u_1, u_2, u_3 about
- Axes g_1, g_2, g_3 , respectively.
- Still missing, angles α_1 between g_1 and g_2 and α_2 between g_2 and g_3 are easy to visualize here.


Split Triad and Three Circles



- Note three circles l, p', k


- Note three circles l, p', k
- Whose centres and radii

Split Triad and Three Circles




- Note three circles l, p', k
- Whose centres and radii
- Squared can be written

Split Triad and Three Circles




- Note three circles l, p', k
- Whose centres and radii
- Squared can be written
- By inspection.

Split Triad and Three Circles




- Note three circles l, p', k
- Whose centres and radii
- Squared can be written
- By inspection.
- $l : \{L(l_1, l_2, l_3), R_L\}$

Split Triad and Three Circles




- Note three circles l, p', k
- Whose centres and radii
- Squared can be written
- By inspection.
- $l : \{L(l_1, l_2, l_3), R_L\}$
- $= \{(-a_1, w_3 \sin \alpha_1,$

Split Triad and Three Circles



- Note three circles l, p', k
- Whose centres and radii
- Squared can be written
- By inspection.
- $l : \{L(l_1, l_2, l_3), R_L\}$
- $= \{(-a_1, w_3 \sin \alpha_1,$
- $w_3 \cos \alpha_1), w_1^2 + w_2^2\}$


Split Triad and Three Circles



- Note three circles l, p', k
- Whose centres and radii
- Squared can be written
- By inspection.
- $l : \{L(l_1, l_2, l_3), R_L\}$
- $= \{(-a_1, w_3 \sin \alpha_1,$
- $w_3 \cos \alpha_1), w_1^2 + w_2^2\}$

- $p' : \{P'(p'_1, p'_2, p'_3), R_P\} = \{(0, 0, d_2 + b \cos \alpha_2), a_2^2 + b^2 \sin^2 \alpha_2\}$


Split Triad and Three Circles



- Note three circles l, p', k
- Whose centres and radii
- Squared can be written
- By inspection.
- $l : \{L(l_1, l_2, l_3), R_L\}$
- $= \{(-a_1, w_3 \sin \alpha_1,$
- $w_3 \cos \alpha_1), w_1^2 + w_2^2\}$

- $p' : \{P'(p'_1, p'_2, p'_3), R_P\} = \{(0, 0, d_2 + b \cos \alpha_2), a_2^2 + b^2 \sin^2 \alpha_2\}$
- $k : \{K(k_1, k_2, k_3), R_K\} = \{(a_2, -b \sin \alpha_2, d_2 + b \cos \alpha_2), s^2\}$

3 Joints Angles and 3 Circular Arcs to a Solution



- Excursion on surface Φ meridian due to rotation u_3 followed by rotation u_2 along latitude circle places W at one of four desired locations on circle I at W_2 .

$I(u_1)$ and $\Phi(u_2, u_3)$ Yields $W(x, y, z)$

$$\bullet I(u_1) : \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -a_1 \\ w_3 \sin \alpha_1 \\ w_3 \cos \alpha_1 \end{bmatrix} + \cos u_1 \begin{bmatrix} w_1 \\ w_2 \cos \alpha_1 \\ -w_2 \sin \alpha_1 \end{bmatrix} +$$

$$\sin u_1 \begin{bmatrix} w_2 \\ -w_1 \cos \alpha_1 \\ w_1 \sin \alpha_1 \end{bmatrix}, \quad \Phi(u_2, u_3) : \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$I(u_1)$ and $\Phi(u_2, u_3)$ Yields $W(x, y, z)$

$$\begin{aligned}
 \bullet \quad I(u_1) : \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \begin{bmatrix} -a_1 \\ w_3 \sin \alpha_1 \\ w_3 \cos \alpha_1 \end{bmatrix} + \cos u_1 \begin{bmatrix} w_1 \\ w_2 \cos \alpha_1 \\ -w_2 \sin \alpha_1 \end{bmatrix} + \\
 &\sin u_1 \begin{bmatrix} w_2 \\ -w_1 \cos \alpha_1 \\ w_1 \sin \alpha_1 \end{bmatrix}, \quad \Phi(u_2, u_3) : \begin{bmatrix} x \\ y \\ z \end{bmatrix} \\
 \bullet &= \begin{bmatrix} \cos u_2(a_2 + s \cos u_3) - \sin u_2(-b \sin \alpha_2 + s \cos \alpha_2 \sin u_3) \\ \sin u_2(a_2 + s \cos u_3) + \cos u_2(-b \sin \alpha_2 + s \cos \alpha_2 \sin u_3) \\ d_2 + b \cos \alpha_2 + s \sin \alpha_2 \sin u_3 \end{bmatrix}
 \end{aligned}$$

$I(u_1)$ and $\Phi(u_2, u_3)$ Yields $W(x, y, z)$

$$\begin{aligned}
 \bullet \quad I(u_1) : \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \begin{bmatrix} -a_1 \\ w_3 \sin \alpha_1 \\ w_3 \cos \alpha_1 \end{bmatrix} + \cos u_1 \begin{bmatrix} w_1 \\ w_2 \cos \alpha_1 \\ -w_2 \sin \alpha_1 \end{bmatrix} + \\
 &\sin u_1 \begin{bmatrix} w_2 \\ -w_1 \cos \alpha_1 \\ w_1 \sin \alpha_1 \end{bmatrix}, \quad \Phi(u_2, u_3) : \begin{bmatrix} x \\ y \\ z \end{bmatrix} \\
 \bullet \quad &= \begin{bmatrix} \cos u_2(a_2 + s \cos u_3) - \sin u_2(-b \sin \alpha_2 + s \cos \alpha_2 \sin u_3) \\ \sin u_2(a_2 + s \cos u_3) + \cos u_2(-b \sin \alpha_2 + s \cos \alpha_2 \sin u_3) \\ d_2 + b \cos \alpha_2 + s \sin \alpha_2 \sin u_3 \end{bmatrix} \\
 \bullet \quad x^2 + y^2 &= (a_2 + s \cos u_3)^2 + (-b \sin \alpha_2 + s \cos \alpha_2 \sin u_3)^2, \\
 \sin u_3 &= \frac{z - d_2 - b \cos \alpha_2}{s \sin \alpha_2}
 \end{aligned}$$

$I(u_1)$ and $\Phi(u_2, u_3)$ Yields $W(x, y, z)$

- $I(u_1) : \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -a_1 \\ w_3 \sin \alpha_1 \\ w_3 \cos \alpha_1 \end{bmatrix} + \cos u_1 \begin{bmatrix} w_1 \\ w_2 \cos \alpha_1 \\ -w_2 \sin \alpha_1 \end{bmatrix} + \sin u_1 \begin{bmatrix} w_2 \\ -w_1 \cos \alpha_1 \\ w_1 \sin \alpha_1 \end{bmatrix}, \quad \Phi(u_2, u_3) : \begin{bmatrix} x \\ y \\ z \end{bmatrix}$
- $= \begin{bmatrix} \cos u_2(a_2 + s \cos u_3) - \sin u_2(-b \sin \alpha_2 + s \cos \alpha_2 \sin u_3) \\ \sin u_2(a_2 + s \cos u_3) + \cos u_2(-b \sin \alpha_2 + s \cos \alpha_2 \sin u_3) \\ d_2 + b \cos \alpha_2 + s \sin \alpha_2 \sin u_3 \end{bmatrix}$
- $x^2 + y^2 = (a_2 + s \cos u_3)^2 + (-b \sin \alpha_2 + s \cos \alpha_2 \sin u_3)^2,$
 $\sin u_3 = \frac{z - d_2 - b \cos \alpha_2}{s \sin \alpha_2}$
- $[(x^2 + y^2 + z^2 - 2d_2z - a_2^2 + d_2^2 - b^2 - s^2) \sin \alpha_2]^2$
 $+ [(z - d_2 - b \cos \alpha_2)^2 - s^2 \sin^2 \alpha_2] 4a_2^2 = 0 \leftarrow \Phi(x, y, z)$

Eliminating u_2, u_3 Produces Quartic Univariate in τ

$$\bullet [1, \cos u_1, \sin u_1] \mathbf{C} \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} = 0, \quad \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} \equiv \begin{bmatrix} 1 + \tau^2 \\ 1 - \tau^2 \\ 2\tau \end{bmatrix}$$

Eliminating u_2, u_3 Produces Quartic Univariate in τ

- $[1, \cos u_1, \sin u_1] \mathbf{C} \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} = 0, \quad \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} \equiv \begin{bmatrix} 1 + \tau^2 \\ 1 - \tau^2 \\ 2\tau \end{bmatrix}$
- $c_{ij} \in \mathbf{C}, \quad c_{ij} = c_{ji}, \quad c_{11} = e^2 + 4fa_2^2, \quad c_{12} = 2(eg - 2ma_2^2 w_2 \sin \alpha_1)$

Eliminating u_2, u_3 Produces Quartic Univariate in τ

- $[1, \cos u_1, \sin u_1] \mathbf{C} \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} = 0, \quad \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} \equiv \begin{bmatrix} 1 + \tau^2 \\ 1 - \tau^2 \\ 2\tau \end{bmatrix}$
- $c_{ij} \in \mathbf{C}, \quad c_{ij} = c_{ji}, \quad c_{11} = e^2 + 4fa_2^2, \quad c_{12} = 2(eg - 2ma_2^2 w_2 \sin \alpha_1)$
- $c_{13} = 2(2ma_2^2 w_1 \sin \alpha_1 - eh), \quad c_{22} = 4(g^2 \sin^2 \alpha_2 + a_2^2 w_2^2 \sin^2 \alpha_1)$

Eliminating u_2, u_3 Produces Quartic Univariate in τ

- $[1, \cos u_1, \sin u_1] \mathbf{C} \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} = 0, \quad \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} \equiv \begin{bmatrix} 1 + \tau^2 \\ 1 - \tau^2 \\ 2\tau \end{bmatrix}$
- $c_{ij} \in \mathbf{C}, \quad c_{ij} = c_{ji}, \quad c_{11} = e^2 + 4fa_2^2, \quad c_{12} = 2(eg - 2ma_2^2 w_2 \sin \alpha_1)$
- $c_{13} = 2(2ma_2^2 w_1 \sin \alpha_1 - eh), \quad c_{22} = 4(g^2 \sin^2 \alpha_2 + a_2^2 w_2^2 \sin^2 \alpha_1)$
- $c_{23} = -4(gh \sin^2 \alpha_2 + a_2^2 w_1 w_2 \sin^2 \alpha_1)$

Eliminating u_2, u_3 Produces Quartic Univariate in τ

- $[1, \cos u_1, \sin u_1] \mathbf{C} \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} = 0, \quad \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} \equiv \begin{bmatrix} 1 + \tau^2 \\ 1 - \tau^2 \\ 2\tau \end{bmatrix}$
- $c_{ij} \in \mathbf{C}, \quad c_{ij} = c_{ji}, \quad c_{11} = e^2 + 4fa_2^2, \quad c_{12} = 2(eg - 2ma_2^2 w_2 \sin \alpha_1)$
- $c_{13} = 2(2ma_2^2 w_1 \sin \alpha_1 - eh), \quad c_{22} = 4(g^2 \sin^2 \alpha_2 + a_2^2 w_2^2 \sin^2 \alpha_1)$
- $c_{23} = -4(gh \sin^2 \alpha_2 + a_2^2 w_1 w_2 \sin^2 \alpha_1)$
- $c_{33} = 4(h^2 \sin^2 \alpha_2 + a_2^2 w_1^2 \sin^2 \alpha_1)$

Eliminating u_2, u_3 Produces Quartic Univariate in τ

- $[1, \cos u_1, \sin u_1] \mathbf{C} \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} = 0, \quad \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} \equiv \begin{bmatrix} 1 + \tau^2 \\ 1 - \tau^2 \\ 2\tau \end{bmatrix}$
- $c_{ij} \in \mathbf{C}, \quad c_{ij} = c_{ji}, \quad c_{11} = e^2 + 4fa_2^2, \quad c_{12} = 2(eg - 2ma_2^2 w_2 \sin \alpha_1)$
- $c_{13} = 2(2ma_2^2 w_1 \sin \alpha_1 - eh), \quad c_{22} = 4(g^2 \sin^2 \alpha_2 + a_2^2 w_2^2 \sin^2 \alpha_1)$
- $c_{23} = -4(gh \sin^2 \alpha_2 + a_2^2 w_1 w_2 \sin^2 \alpha_1)$
- $c_{33} = 4(h^2 \sin^2 \alpha_2 + a_2^2 w_1^2 \sin^2 \alpha_1)$
- $e = a_1^2 - a_2^2 + d_2^2 - b^2 - s^2 + w_1^2 + w_2^2 + w_3^2 - 2d_2 w_3 \cos \alpha_1$

Eliminating u_2, u_3 Produces Quartic Univariate in τ

- $[1, \cos u_1, \sin u_1] \mathbf{C} \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} = 0, \quad \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} \equiv \begin{bmatrix} 1 + \tau^2 \\ 1 - \tau^2 \\ 2\tau \end{bmatrix}$
- $c_{ij} \in \mathbf{C}, \quad c_{ij} = c_{ji}, \quad c_{11} = e^2 + 4fa_2^2, \quad c_{12} = 2(eg - 2ma_2^2 w_2 \sin \alpha_1)$
- $c_{13} = 2(2ma_2^2 w_1 \sin \alpha_1 - eh), \quad c_{22} = 4(g^2 \sin^2 \alpha_2 + a_2^2 w_2^2 \sin^2 \alpha_1)$
- $c_{23} = -4(gh \sin^2 \alpha_2 + a_2^2 w_1 w_2 \sin^2 \alpha_1)$
- $c_{33} = 4(h^2 \sin^2 \alpha_2 + a_2^2 w_1^2 \sin^2 \alpha_1)$
- $e = a_1^2 - a_2^2 + d_2^2 - b^2 - s^2 + w_1^2 + w_2^2 + w_3^2 - 2d_2 w_3 \cos \alpha_1$
- $f = (w_3 \cos \alpha_1 - d_2 - b \cos \alpha_2)^2 - s^2 \sin^2 \alpha_2$

Eliminating u_2, u_3 Produces Quartic Univariate in τ

- $[1, \cos u_1, \sin u_1] \mathbf{C} \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} = 0, \quad \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} \equiv \begin{bmatrix} 1 + \tau^2 \\ 1 - \tau^2 \\ 2\tau \end{bmatrix}$
- $c_{ij} \in \mathbf{C}, \quad c_{ij} = c_{ji}, \quad c_{11} = e^2 + 4fa_2^2, \quad c_{12} = 2(eg - 2ma_2^2 w_2 \sin \alpha_1)$
- $c_{13} = 2(2ma_2^2 w_1 \sin \alpha_1 - eh), \quad c_{22} = 4(g^2 \sin^2 \alpha_2 + a_2^2 w_2^2 \sin^2 \alpha_1)$
- $c_{23} = -4(gh \sin^2 \alpha_2 + a_2^2 w_1 w_2 \sin^2 \alpha_1)$
- $c_{33} = 4(h^2 \sin^2 \alpha_2 + a_2^2 w_1^2 \sin^2 \alpha_1)$
- $e = a_1^2 - a_2^2 + d_2^2 - b^2 - s^2 + w_1^2 + w_2^2 + w_3^2 - 2d_2 w_3 \cos \alpha_1$
- $f = (w_3 \cos \alpha_1 - d_2 - b \cos \alpha_2)^2 - s^2 \sin^2 \alpha_2$
- $g = d_2 w_2 \sin \alpha_1 - a_1 w_1, \quad h = d_2 w_1 \sin \alpha_1 + a_1 w_1$

Eliminating u_2, u_3 Produces Quartic Univariate in τ

- $[1, \cos u_1, \sin u_1] \mathbf{C} \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} = 0, \quad \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} \equiv \begin{bmatrix} 1 + \tau^2 \\ 1 - \tau^2 \\ 2\tau \end{bmatrix}$
- $c_{ij} \in \mathbf{C}, \quad c_{ij} = c_{ji}, \quad c_{11} = e^2 + 4fa_2^2, \quad c_{12} = 2(eg - 2ma_2^2 w_2 \sin \alpha_1)$
- $c_{13} = 2(2ma_2^2 w_1 \sin \alpha_1 - eh), \quad c_{22} = 4(g^2 \sin^2 \alpha_2 + a_2^2 w_2^2 \sin^2 \alpha_1)$
- $c_{23} = -4(gh \sin^2 \alpha_2 + a_2^2 w_1 w_2 \sin^2 \alpha_1)$
- $c_{33} = 4(h^2 \sin^2 \alpha_2 + a_2^2 w_1^2 \sin^2 \alpha_1)$
- $e = a_1^2 - a_2^2 + d_2^2 - b^2 - s^2 + w_1^2 + w_2^2 + w_3^2 - 2d_2 w_3 \cos \alpha_1$
- $f = (w_3 \cos \alpha_1 - d_2 - b \cos \alpha_2)^2 - s^2 \sin^2 \alpha_2$
- $g = d_2 w_2 \sin \alpha_1 - a_1 w_1, \quad h = d_2 w_1 \sin \alpha_1 + a_1 w_1$
- $m = w_3 \cos \alpha_1 - d_2 - b \cos \alpha_2$

Eliminating u_2, u_3 Produces Quartic Univariate in τ

- $[1, \cos u_1, \sin u_1] \mathbf{C} \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} = 0, \quad \begin{bmatrix} 1 \\ \cos u_1 \\ \sin u_1 \end{bmatrix} \equiv \begin{bmatrix} 1 + \tau^2 \\ 1 - \tau^2 \\ 2\tau \end{bmatrix}$
- $c_{ij} \in \mathbf{C}, \quad c_{ij} = c_{ji}, \quad c_{11} = e^2 + 4fa_2^2, \quad c_{12} = 2(eg - 2ma_2^2 w_2 \sin \alpha_1)$
- $c_{13} = 2(2ma_2^2 w_1 \sin \alpha_1 - eh), \quad c_{22} = 4(g^2 \sin^2 \alpha_2 + a_2^2 w_2^2 \sin^2 \alpha_1)$
- $c_{23} = -4(gh \sin^2 \alpha_2 + a_2^2 w_1 w_2 \sin^2 \alpha_1)$
- $c_{33} = 4(h^2 \sin^2 \alpha_2 + a_2^2 w_1^2 \sin^2 \alpha_1)$
- $e = a_1^2 - a_2^2 + d_2^2 - b^2 - s^2 + w_1^2 + w_2^2 + w_3^2 - 2d_2 w_3 \cos \alpha_1$
- $f = (w_3 \cos \alpha_1 - d_2 - b \cos \alpha_2)^2 - s^2 \sin^2 \alpha_2$
- $g = d_2 w_2 \sin \alpha_1 - a_1 w_1, \quad h = d_2 w_1 \sin \alpha_1 + a_1 w_1$
- $m = w_3 \cos \alpha_1 - d_2 - b \cos \alpha_2$
- $\tau \rightarrow \cos u_1, \sin u_1, \text{ up to four real solutions.}$

Computing u_3 , u_2

- From parametrization of $\Phi(u_2, u_3) \rightarrow z$ recall

$$\sin u_3 = \frac{z - d_2 - b \cos \alpha_2}{s \sin \alpha_2}.$$

Computing u_3 , u_2

- From parametrization of $\Phi(u_2, u_3) \rightarrow z$ recall

$$\sin u_3 = \frac{z - d_2 - b \cos \alpha_2}{s \sin \alpha_2}.$$
- So $\cos u_3 = \pm \sqrt{1 - \sin^2 u_3}.$

Computing u_3 , u_2

- From parametrization of $\Phi(u_2, u_3) \rightarrow z$ recall $\sin u_3 = \frac{z - d_2 - b \cos \alpha_2}{s \sin \alpha_2}$.
- So $\cos u_3 = \pm \sqrt{1 - \sin^2 u_3}$.
- To check \pm use $p + q - x^2 - y^2 = 0$

Computing u_3 , u_2

- From parametrization of $\Phi(u_2, u_3) \rightarrow z$ recall $\sin u_3 = \frac{z - d_2 - b \cos \alpha_2}{s \sin \alpha_2}$.
- So $\cos u_3 = \pm \sqrt{1 - \sin^2 u_3}$.
- To check \pm use $p + q - x^2 - y^2 = 0$
- Where $p = (a_2 + s \cos u_3)^2$ and $q = (s \cos \alpha_2 \sin u_3 - b \sin \alpha_2)^2$.



Computing u_3 , u_2

- From parametrization of $\Phi(u_2, u_3) \rightarrow z$ recall $\sin u_3 = \frac{z - d_2 - b \cos \alpha_2}{s \sin \alpha_2}$.
- So $\cos u_3 = \pm \sqrt{1 - \sin^2 u_3}$.
- To check \pm use $p + q - x^2 - y^2 = 0$
- Where $p = (a_2 + s \cos u_3)^2$ and $q = (s \cos \alpha_2 \sin u_3 - b \sin \alpha_2)^2$.
- From parametrization of $\Phi(u_2, u_3) \rightarrow x, y$ recall

Computing u_3 , u_2

- From parametrization of $\Phi(u_2, u_3) \rightarrow z$ recall $\sin u_3 = \frac{z - d_2 - b \cos \alpha_2}{s \sin \alpha_2}$.
- So $\cos u_3 = \pm \sqrt{1 - \sin^2 u_3}$.
- To check \pm use $p + q - x^2 - y^2 = 0$
- Where $p = (a_2 + s \cos u_3)^2$ and $q = (s \cos \alpha_2 \sin u_3 - b \sin \alpha_2)^2$.
- From parametrization of $\Phi(u_2, u_3) \rightarrow x, y$ recall
- $\cos u_2 = \frac{px + qy}{x^2 + y^2}$, $\sin u_2 = \frac{py - qx}{x^2 + y^2}$.

Summing Up: 4 Solutions and 4 Postures



Three Surfaces

- Recall the cyclid:

$$C_\infty(C_\infty + c_{Cz}z + c_C) + c_{z2}z^2 + c_zz + c = 0.$$

Three Surfaces

- Recall the cyclid:

$$C_\infty(C_\infty + c_{Cz}z + c_C) + c_{z2}z^2 + c_zz + c = 0.$$

- Consider a sphere centred on L , radius squared R :

$$C_\infty - k_x x - k_y y - k_z z - k = 0$$

Three Surfaces

- Recall the cyclid:

$$C_\infty(C_\infty + c_{Cz}z + c_C) + c_{z2}z^2 + c_zz + c = 0.$$

- Consider a sphere centred on L , radius squared R :

$$C_\infty - k_x x - k_y y - k_z z - k = 0$$

- And a plane e on L and normal to g_1 :

$$e_x x + e_y y + e_z z + e = 0.$$

Three Surfaces

- Recall the cyclid:

$$C_\infty(C_\infty + c_{Cz}z + c_C) + c_{z2}z^2 + c_zz + c = 0.$$

- Consider a sphere centred on L , radius squared R :

$$C_\infty - k_x x - k_y y - k_z z - k = 0$$

- And a plane e on L and normal to g_1 :

$$e_x x + e_y y + e_z z + e = 0.$$

- Replace C_∞ in the cyclid with linear form in sphere equation.

Three Surfaces

- Recall the cyclid:

$$C_\infty(C_\infty + c_{Cz}z + c_C) + c_{z2}z^2 + c_zz + c = 0.$$

- Consider a sphere centred on L , radius squared R :

$$C_\infty - k_x x - k_y y - k_z z - k = 0$$

- And a plane e on L and normal to g_1 :

$$e_x x + e_y y + e_z z + e = 0.$$

- Replace C_∞ in the cyclid with linear form in sphere equation.

- $(k_x x + k_y y + k_z z + k)[(k_x x + k_y y + k_z z + k) + c_{Cz}z + c_C] + c_{z2}z^2 + c_zz + c = 0$

Three Surfaces

- Recall the cyclid:

$$C_\infty(C_\infty + c_{Cz}z + c_C) + c_{z2}z^2 + c_zz + c = 0.$$

- Consider a sphere centred on L , radius squared R :

$$C_\infty - k_x x - k_y y - k_z z - k = 0$$

- And a plane e on L and normal to g_1 :


$$e_x x + e_y y + e_z z + e = 0.$$

- Replace C_∞ in the cyclid with linear form in sphere equation.

- $(k_x x + k_y y + k_z z + k)[(k_x x + k_y y + k_z z + k) + c_{Cz}z + c_C] + c_{z2}z^2 + c_zz + c = 0$


- Thus producing a quadric surface that yields the four values of S_i when combined with the sphere and plane.

All Four Surfaces Together



- Eliminating the absolute conic $C_\infty = x^2 + y^2 + z^2$ from the cyclid with the linear form from the sphere produces a cylinder with an elliptic right section.

All Four Surfaces Together



- Eliminating the absolute conic $C_\infty = x^2 + y^2 + z^2$ from the cyclid with the linear form from the sphere produces a cylinder with an elliptic right section.
- Intersection of the sphere, cylinder and plane yields the four required points $S_i, i = 1, 2, 3, 4$.

What Was Accomplished

- Choosing the second revolute axis g_2 as the reference frame split the problem in two, nicely.

What Was Accomplished

- Choosing the second revolute axis g_2 as the reference frame split the problem in two, nicely.
- A fourth order surface, the cyclid, is generated, naturally, by sweeping the wrist point W , initially in home position, about axes $g_{1,2}$

What Was Accomplished

- Choosing the second revolute axis g_2 as the reference frame split the problem in two, nicely.
- A fourth order surface, the cyclid, is generated, naturally, by sweeping the wrist point W , initially in home position, about axes $g_{1,2}$
- Thus parameterizing it in terms of joint angles $u_{2,3}$.

What Was Accomplished

- Choosing the second revolute axis g_2 as the reference frame split the problem in two, nicely.
- A fourth order surface, the cyclid, is generated, naturally, by sweeping the wrist point W , initially in home position, about axes $g_{1,2}$
- Thus parameterizing it in terms of joint angles $u_{2,3}$.
- A circle, in terms of joint angle u_1 , is generated by revolving the given wrist centre W_0 about axis g_1 .

What Was Accomplished

- Choosing the second revolute axis g_2 as the reference frame split the problem in two, nicely.
- A fourth order surface, the cyclid, is generated, naturally, by sweeping the wrist point W , initially in home position, about axes $g_{1,2}$
- Thus parameterizing it in terms of joint angles $u_{2,3}$.
- A circle, in terms of joint angle u_1 , is generated by revolving the given wrist centre W_0 about axis g_1 .
- Separating joint variables in this way renders the coefficients in the equation set quite compact

What Was Accomplished

- Choosing the second revolute axis g_2 as the reference frame split the problem in two, nicely.
- A fourth order surface, the cyclid, is generated, naturally, by sweeping the wrist point W , initially in home position, about axes $g_{1,2}$
- Thus parameterizing it in terms of joint angles $u_{2,3}$.
- A circle, in terms of joint angle u_1 , is generated by revolving the given wrist centre W_0 about axis g_1 .
- Separating joint variables in this way renders the coefficients in the equation set quite compact
- And Identifies the cyclid as the invariant or architectural geometric element; the circle is the variable element.

Questions to Be Investigated

- If the wrist point tracks a straight line what is the nature of the curves it traces on the cyclid? *E.g.*, are they geodesics? No, they are not

Questions to Be Investigated

- If the wrist point tracks a straight line what is the nature of the curves it traces on the cyclid? *E.g.*, are they geodesics? No, they are not
- Because all straight lines, generally skew to axis g_1 , will generate one-sheet hyperboloids of revolution to intersect the cyclid in a curve of eighth degree. Can it be reduced?

Questions to Be Investigated



- If the wrist point tracks a straight line what is the nature of the curves it traces on the cyclid? *E.g.*, are they geodesics? No, they are not
- Because all straight lines, generally skew to axis g_1 , will generate one-sheet hyperboloids of revolution to intersect the cyclid in a curve of eighth degree. Can it be reduced?
- If the wrist point tracks a circle the surface of revolution about g_1 will be a cyclid, similar to the one generated by W when swept by revolutions about g_2 and g_3 . What is the intersection curve like in this case?

Questions to Be Investigated

- If the wrist point tracks a straight line what is the nature of the curves it traces on the cyclid? *E.g.*, are they geodesics? No, they are not
- Because all straight lines, generally skew to axis g_1 , will generate one-sheet hyperboloids of revolution to intersect the cyclid in a curve of eighth degree. Can it be reduced?
- If the wrist point tracks a circle the surface of revolution about g_1 will be a cyclid, similar to the one generated by W when swept by revolutions about g_2 and g_3 . What is the intersection curve like in this case?
- How may the cyclid-circle model be exploited to gain insight in velocity analysis? ... acceleration ... ?



Two Special Cases

- Recall the cyclid swept under the auspices of revolution of W about $g_{2,3}$ through full circles $u_{2,3} = 2\pi$.





Two Special Cases

- Recall the cyclid swept under the auspices of revolution of W about $g_{2,3}$ through full circles $u_{2,3} = 2\pi$.
- This becomes an elliptical cylinder when combined with a sphere that contains the circle swept by W_0 as it rotates about axis g_1 through $u_1 = 2\pi$.





Two Special Cases

- Recall the cyclid swept under the auspices of revolution of W about $g_{2,3}$ through full circles $u_{2,3} = 2\pi$.
- This becomes an elliptical cylinder when combined with a sphere that contains the circle swept by W_0 as it rotates about axis g_1 through $u_1 = 2\pi$.
- A case where $\exists g_2 \cap g_3 \equiv a_2 = 0$




Two Special Cases

- Recall the cyclid swept under the auspices of revolution of W about $g_{2,3}$ through full circles $u_{2,3} = 2\pi$.
- This becomes an elliptical cylinder when combined with a sphere that contains the circle swept by W_0 as it rotates about axis g_1 through $u_1 = 2\pi$.
- A case where $\exists g_2 \cap g_3 \equiv a_2 = 0$
- And another where $\alpha_2 = 0$, $a_2 \neq 0$ will be considered.




Axes g_2 and g_3 Intersect

- Surface is a bi-truncated sphere on $P = Q$, radius $+\sqrt{b^2 + s^2}$.




Axes g_2 and g_3 Intersect

- Surface is a bi-truncated sphere on $P = Q$, radius $+\sqrt{b^2 + s^2}$.
- Cyclid equation degenerates, with $a_2 = 0$, to $C_\infty - 2d_2z + d^2 - b^2 - s^2 = 0$.




Axes g_2 and g_3 Intersect

- Surface is a bi-truncated sphere on $P = Q$, radius $+\sqrt{b^2 + s^2}$.
- Cyclid equation degenerates, with $a_2 = 0$, to $C_\infty - 2d_2z + d^2 - b^2 - s^2 = 0$.
- Region lies between $z = d_2b \cos \alpha_2 \pm s \sin \alpha_2$.




Axes g_2 and g_3 Intersect

- Surface is a bi-truncated sphere on $P = Q$, radius $+\sqrt{b^2 + s^2}$.
- Cyclid equation degenerates, with $a_2 = 0$, to $C_\infty - 2d_2z + d^2 - b^2 - s^2 = 0$.
- Region lies between $z = d_2b \cos \alpha_2 \pm s \sin \alpha_2$.
- Circle on axis g_1 intersects it in two places




Axes g_2 and g_3 Intersect

- Surface is a bi-truncated sphere on $P = Q$, radius $+\sqrt{b^2 + s^2}$.
- Cyclid equation degenerates, with $a_2 = 0$, to $C_\infty - 2d_2z + d^2 - b^2 - s^2 = 0$.
- Region lies between $z = d_2b \cos \alpha_2 \pm s \sin \alpha_2$.
- Circle on axis g_1 intersects it in two places
- But sign test for $\cos u_3 = \pm \sqrt{1 - \sin^2 u_2}$ fails




Axes g_2 and g_3 Intersect

- Surface is a bi-truncated sphere on $P = Q$, radius $+\sqrt{b^2 + s^2}$.
- Cyclid equation degenerates, with $a_2 = 0$, to $C_\infty - 2d_2z + d^2 - b^2 - s^2 = 0$.
- Region lies between $z = d_2b \cos \alpha_2 \pm s \sin \alpha_2$.
- Circle on axis g_1 intersects it in two places
- But sign test for $\cos u_3 = \pm \sqrt{1 - \sin^2 u_2}$ fails
- Because $p = (a_2 + s \cos u_3)^2$ returns a unique value



Axes g_2 and g_3 Intersect

- Surface is a bi-truncated sphere on $P = Q$, radius $+\sqrt{b^2 + s^2}$.
- Cyclid equation degenerates, with $a_2 = 0$, to $C_\infty - 2d_2z + d^2 - b^2 - s^2 = 0$.
- Region lies between $z = d_2b \cos \alpha_2 \pm s \sin \alpha_2$.
- Circle on axis g_1 intersects it in two places
- But sign test for $\cos u_3 = \pm \sqrt{1 - \sin^2 u_2}$ fails
- Because $p = (a_2 + s \cos u_3)^2$ returns a unique value
- So the solution admits two valid values as $u_3, u_3 + \pi$



$$g_2 \parallel g_3, g_1 \cap g_2, g_1 \parallel g_2$$

- Since $\sin \alpha_2 = 0$ surface is region in either plane $z = d_2 \pm b$.

$$g_2 \parallel g_3, g_1 \cap g_2, g_1 \parallel g_2$$

- Since $\sin \alpha_2 = 0$ surface is region in either plane $z = d_2 \pm b$.
- It's an annulus and \pm denotes $\alpha_2 = 0, \pi$, respectively.

$$g_2 \parallel g_3, g_1 \cap g_2, g_1 \parallel g_2$$

- Since $\sin \alpha_2 = 0$ surface is region in either plane $z = d_2 \pm b$.
- It's an annulus and \pm denotes $\alpha_2 = 0, \pi$, respectively.
- Interior/exterior radii are $a_2 \pm s$.

$$g_2 \parallel g_3, g_1 \cap g_2, g_1 \parallel g_2$$

- Since $\sin \alpha_2 = 0$ surface is region in either plane $z = d_2 \pm b$.
- It's an annulus and \pm denotes $\alpha_2 = 0, \pi$, respectively.
- Interior/exterior radii are $a_2 \pm s$.
- Circle $x, y, (z)$ equation triad gives

$$d_2 \pm b - w_3 \cos \alpha_1 + \sin \alpha_1 (w_2 \cos u_1 - w_1 \sin u_1) = 0.$$

$$g_2 \parallel g_3, g_1 \cap g_2, g_1 \parallel g_2$$

- Since $\sin \alpha_2 = 0$ surface is region in either plane $z = d_2 \pm b$.
- It's an annulus and \pm denotes $\alpha_2 = 0, \pi$, respectively.
- Interior/exterior radii are $a_2 \pm s$.
- Circle $x, y, (z)$ equation triad gives
 $d_2 \pm b - w_3 \cos \alpha_1 + \sin \alpha_1 (w_2 \cos u_1 - w_1 \sin u_1) = 0$.
- Maybe two real solutions after τ substitution; circle \cap plane

$$g_2 \parallel g_3, g_1 \cap g_2, g_1 \parallel g_2$$

- Since $\sin \alpha_2 = 0$ surface is region in either plane $z = d_2 \pm b$.
- It's an annulus and \pm denotes $\alpha_2 = 0, \pi$, respectively.
- Interior/exterior radii are $a_2 \pm s$.
- Circle $x, y, (z)$ equation triad gives

$$d_2 \pm b - w_3 \cos \alpha_1 + \sin \alpha_1 (w_2 \cos u_1 - w_1 \sin u_1) = 0.$$
- Maybe two real solutions after τ substitution; circle \cap plane
- Subject to $a_2 - s \leq \sqrt{x^2 + y^2} \leq a_2 + s$.

$$g_2 \parallel g_3, g_1 \cap g_2, g_1 \parallel g_2$$

- Since $\sin \alpha_2 = 0$ surface is region in either plane $z = d_2 \pm b$.
- It's an annulus and \pm denotes $\alpha_2 = 0, \pi$, respectively.
- Interior/exterior radii are $a_2 \pm s$.
- Circle $x, y, (z)$ equation triad gives

$$d_2 \pm b - w_3 \cos \alpha_1 + \sin \alpha_1 (w_2 \cos u_1 - w_1 \sin u_1) = 0.$$
- Maybe two real solutions after τ substitution; circle \cap plane
- Subject to $a_2 - s \leq \sqrt{x^2 + y^2} \leq a_2 + s$.
- Use surface equation $(x, y), z$ triad $x = s \cos(u_2 \pm u_3)$

$$+ a_2 \cos u_2, y = s \sin(u_2 \pm u_3) + a_2 \sin u_2 \rightarrow \cos u_3 = \frac{x^2 + y^2 - a_2^2 - s^2}{2a_2 s}.$$

$$g_2 \parallel g_3, g_1 \cap g_2, g_1 \parallel g_2$$

- Since $\sin \alpha_2 = 0$ surface is region in either plane $z = d_2 \pm b$.
- It's an annulus and \pm denotes $\alpha_2 = 0, \pi$, respectively.
- Interior/exterior radii are $a_2 \pm s$.
- Circle $x, y, (z)$ equation triad gives

$$d_2 \pm b - w_3 \cos \alpha_1 + \sin \alpha_1 (w_2 \cos u_1 - w_1 \sin u_1) = 0.$$
- Maybe two real solutions after τ substitution; circle \cap plane
- Subject to $a_2 - s \leq \sqrt{x^2 + y^2} \leq a_2 + s$.
- Use surface equation $(x, y), z$ triad $x = s \cos(u_2 \pm u_3)$
 $+ a_2 \cos u_2, y = s \sin(u_2 \pm u_3) + a_2 \sin u_2 \rightarrow \cos u_3 = \frac{x^2 + y^2 - a_2^2 - s^2}{2a_2 s}.$
- For each u_1 , two u_3 ; $u_3 + u'_3 = 2\pi$.

$g_2 \parallel g_3, g_1 \cap g_2, g_1 \parallel g_2$

- Since $\sin \alpha_2 = 0$ surface is region in either plane $z = d_2 \pm b$.
- It's an annulus and \pm denotes $\alpha_2 = 0, \pi$, respectively.
- Interior/exterior radii are $a_2 \pm s$.
- Circle $x, y, (z)$ equation triad gives

$$d_2 \pm b - w_3 \cos \alpha_1 + \sin \alpha_1 (w_2 \cos u_1 - w_1 \sin u_1) = 0.$$
- Maybe two real solutions after τ substitution; circle \cap plane
- Subject to $a_2 - s \leq \sqrt{x^2 + y^2} \leq a_2 + s$.
- Use surface equation $(x, y), z$ triad $x = s \cos(u_2 \pm u_3)$
 $+ a_2 \cos u_2, y = s \sin(u_2 \pm u_3) + a_2 \sin u_2 \rightarrow \cos u_3 = \frac{x^2 + y^2 - a_2^2 - s^2}{2a_2 s}.$
- For each u_1 , two u_3 ; $u_3 + u'_3 = 2\pi$.
- If $\exists g_1 \cap g_2$, reverse rôles of circles k and l .